


JCO7 Rec'd PCT/PTO 27 DEC 2001

FORM PTO-1390 (REV. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 442P090
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 10/030294
INTERNATIONAL APPLICATION NO. PCT/JP00/04261	INTERNATIONAL FILING DATE 28 June 2000	PRIORITY DATE CLAIMED 29 June 1999	
TITLE OF INVENTION GENE ENCODING PROMOTER REGION OF TUMOR SUPPRESSOR GENE P51 AND USE THEREOF			
APPLICANT(S) FOR DO/EO/US Toshiyuki Sakai, Shigehide Kagaya, Takamichi Sato, Yoshikazu Sukenaga and Hideji Fujii			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
<ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). <ol style="list-style-type: none"> a. <input checked="" type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 			
Items 11 to 20 below concern document(s) or information included:			
<ol style="list-style-type: none"> 11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. 14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input checked="" type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information: <ol style="list-style-type: none"> 1. Disk containing sequence listing; and 27-Page printed sequence listing. 2. Copy of the International Search Report; 3. Copy of Form PCT/IB/308; and 4. Copy of Form PCT/IB/304. 			

U.S. APPLICATION NO. (if known) 10/030294		INTERNATIONAL APPLICATION NO. PCT/JP00/04261		ATTORNEY'S DOCKET NUMBER 442P090	
21. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. \$1040.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 890.00	
				\$ 0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total claims	15 - 20 =	0	x \$18.00	\$ 0.00	
Independent claims	2 - 3 =	0	x \$84.00	\$ 0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$280.00	\$ 280.00	
TOTAL OF ABOVE CALCULATIONS =				\$ 1,170.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$ 0	
SUBTOTAL =				\$ 1,170.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$ 0	
TOTAL NATIONAL FEE =				\$ 1,170.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$ 40.00	
TOTAL FEES ENCLOSED =				\$ 1,210.00	
				Amount to be refunded:	\$
				charged:	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>1,170.00 & 40.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>14-0930</u> . A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
Kevin S. Lemack Nields & Lemack 176 E. Main Street Westboro, MA 01581			<div style="text-align: center;">  SIGNATURE </div> <div style="text-align: center;"> Kevin S. Lemack NAME </div> <div style="text-align: center;"> 32,579 REGISTRATION NUMBER </div>		

3/prb

1

DESCRIPTION

GENE ENCODING PROMOTER REGION OF TUMOR SUPPRESSOR GENE

P51 AND USE THEREOF

TECHNICAL FIELD

The present invention relates to a gene encoding the promoter region of a protein p51 which is capable of inducing cell death, suppressing cell growth, etc. and a gene encoding the 5'-untranslated region of p51. The present invention also relates to a series of uses of these genes.

BACKGROUND ART

The tumor suppressor gene p53 encodes a protein having a variety of functions such as the induction of apoptosis, cell cycle arrest, the repairing of DNA damages, and the like. It has also been shown that p53 is a transcription factor which regulates the expression of various proteins.

Currently this protein is believed to play a central role in the control of cell growth. It has also been reported that p53 mutation is observed in about half of tumor cells which is mainly responsible for abnormal growth and resistance against anti-cancer agents.

Although no proteins were known that have a p53-like structure and function until very recently, but two novel p53-like molecules were recently reported. One

is p73 (Cell, 90:809-819, 1997), and the other is p51 (Nature Medicine, 4:839-843, 1998). Concerning p51, a protein encoded by the same gene is reported as p63 (Molecular Cell, 2:305-316, 1998). Structurally p51 is very similar to p53, and like p53 it is capable of activating the transcription of a cell cycle progression suppressor protein p21, suggesting that it is homologous to p53 functionally as well. Furthermore, it is reported that p51, as is p53, is capable of inducing cell death, suppressing cell growth, and the like. It has been shown, however, that p51 is different from p53 in that no mutation in p51 has been observed in tumor cells. It is being elucidated that p51 is expressed in very limited organs such as muscle cells.

DISCLOSURE OF THE INVENTION

It is an object of the present invention to identify the gene encoding the promoter region of a tumor suppressor gene p51 and the gene encoding the 5'-untranslated region of p51, thereby to provide a method of cloning these genes, a method of screening novel drugs using these genes, and a novel method of gene therapy using these genes, and the like.

After intensive and extensive research to find a gene fragment containing the p51 promoter in a human genomic library, the inventor of the present invention has identified a gene sequence having the p51

promoter region, and a gene sequence encoding the 5'-untranslated region of p51, and thereby has completed the present invention.

Thus, a human genome fragment containing a
5 same sequence as cDNA complementary to p51 mRNA was
screened and isolated by the plaque hybridization
screening method. Furthermore, the base sequence of
this gene fragment was determined to confirm that it
contains regions that are estimated to be the p51
10 promoter region and the 5'-untranslated region of p51.
A plasmid was also constructed in which the gene
fragment is ligated to the luciferase reporter gene,
and it was then introduced into a human cell line to
construct a transformant. By further analyzing this
15 transformant, it was found out that said gene can work
as a promoter. Furthermore, there was found a method
of screening an agent that acts on the p51 promoter
using transformant, and a possibility that a pharma-
ceutical agent that enhances this promoter activity
20 acts as a therapeutic agent such as an anti-cancer
agent for diseases in which abnormality in p53-
dependent apoptosis is involved.

Thus, the present invention relates to a gene
encoding the p51 promoter region shown in the following
25 (1), (2), (3), (4), (5), or (6):

(1) DNA that encodes the p51 promoter region
having the base sequence as set forth in SEQ ID NO: 1
of the sequence listing;

(2) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 1 of the sequence listing, and that has p51

5 promoter activity;

(3) DNA that hybridizes to the base sequence as set forth in SEQ ID NO: 1 of the sequence listing under a stringent condition, and that has p51 promoter activity;

10 (4) DNA that has the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that encodes the p51 promoter region and the 5'-untranslated region of p51;

(5) DNA that has a base sequence in which one
15 or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that has p51 promoter activity; and

(6) DNA that hybridizes to the base sequence
20 as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition, and that has p51 promoter activity.

The present invention also relates to a gene encoding the 5'-untranslated region of p51 shown in the
25 following (7), (8), or (9):

(7) DNA that has a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing;

(8) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that has a function similar to that of DNA in the above (7); and

(9) DNA that hybridizes to DNA comprising the base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition, and that has a function similar to that of DNA in the above (7).

The present invention further relates to a recombinant plasmid comprising the gene in the above (1) to (6).

The present invention further relates to a transformant or a transductant comprising the above recombinant plasmid.

The present invention further relates to nucleic acid probes comprising all or parts of the base sequence of the gene in the above (1) to (9).

The present invention further relates to a method of cloning p51 promoter by hybridization using the above nucleic acid probe.

The present invention further relates to a DNA sequence that is an antisense to all or part of the base sequence of the gene in the above (1) to (9), and that can modify the function of p51 promoter activity.

The present invention further relates to a

RNA sequence that is an antisense to all or part of the base sequence of the gene in the above (1) to (9), and that can modify the function of p51 promoter activity.

The present invention further relates to a method of screening drugs that act on the p51 promoter using the above transformant or the above transductant.

The present invention further relates to a compound that enhances or inhibits the expression of p51 gene, said compound being selected by the above screening method.

The present invention further relates to a pharmaceutical formulation comprising the above DNA or the above RNA.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram showing a restriction map of pGL2-neo/p51 promoter. In the figure, amp represents an ampicillin-resistant gene, neo represents a neomycin-resistant gene, and luc represents the luciferase gene. The name of a restriction enzyme that has a * mark on the left means that the vector can be only cleaved at the point.

Figure 2 shows an analysis in which the amount of p51 expressed was analyzed using the technique of Northern blotting after a cell line HCT116 derived from colon cancer, was subjected to various stimulations. In each lane is blotted 20 μ g of RNA from: lane 1, 2: cells treated with taxol (1 μ g/ml, 10

5 $\mu\text{g/ml}$), lane 3, 4: cells treated with trichostatin A
 (0.1 $\mu\text{g/ml}$, 1 $\mu\text{g/ml}$), lane 5, 6: cells treated with
 radiation (50 J/m^2 , 100 J/m^2), lane 7, 8: cells treated
 with vitamin D₃ (10 nM, 100 nM), lane 9, 10: cells
 11, 12: cells treated with wortmannin (0.1 μM , 1 μM).
 A band that reacts with the p51 probe is observed at
 the position indicated by an arrow.

Figure 3 shows a cell line HCT116 derived
 10 from colon cancer, into which 2 μg each of pGL2-neo/p51
 promoter or pGL2-neo vector was introduced and 24 hours
 later, 0 or 1 $\mu\text{g/ml}$ trichostatin A (TsA) was added and
 was further cultured for 24 hours. By determining the
 luciferase activity in these cells, the promoter
 15 activity of the isolated gene fragments was examined.
 As a negative control cell, cells into which 2 μg of
 pCDNA3/lacZ was introduced were used. As a result, it
 was found that an extract of the pGL2-neo/p51 promoter-
 introduced cells exhibits a significantly higher
 20 fluorescence intensity than that of the pGL2-neo-
 introduced cells. The administration of trichostatin A
 further enhanced the chemiluminescence intensity in the
 extract of the pGL2-neo/p51 promoter-introduced cells.

BEST MODE FOR CARRYING OUT THE INVENTION

25 The present invention will now be explained
 in further details below.

(1) Gene having the p51 promoter region of the present

invention

(1-1) In order to isolate a gene having the p51 promoter region of the present invention, a human genomic DNA library is constructed, and from the library, using part of the sequence described in the above-mentioned article in Nature Medicine, a cDNA molecule that hybridizes to a cDNA complementary to the 5'-untranslated region of p51 mRNA can be isolated and cloned resulting in the isolation of a gene encoding the p51 promoter region.

The base sequence of a gene encoding the p51 promoter region of the present invention is as set forth in SEQ ID NO: 1. In this gene sequence, TATA boxes have been observed in the bases in positions from 5630 to 5636 and in positions from 5659 to 5665 of the above same sequence listing. They are located at 47 bases and 18 bases upstream of a gene sequence (GenBank AB016072) reported as the sequence of p51A protein mRNA (p51AmRNA), respectively. Furthermore, the binding site of MEF2 known to be a muscle cell-specific transcription factor is located in the bases at positions from 1211 to 1220 of the same sequence listing. In a search of the database for genes having a homology with the gene encoding The p51 promoter region as set forth in SEQ ID NO: 1 of the sequence listing, no sequences had a homology of 30% or higher over the entire length.

Furthermore, for the purpose of confirming the function of the gene having the base sequence as

set forth in SEQ ID NO: 1 of the sequence listing, as shown in the following Example, a plasmid was constructed in which a gene having the base sequence as set forth in SEQ ID NO: 1 of the sequence listing and the luciferase reporter gene were ligated, which was then expressed in a cultured human cell line to construct a transformant. The transformant significantly expressed the luciferase reporter gene as compared to the control group. Furthermore, a simple screening using the transformant led to the discovery of the p51 promoter activating ability in trichostatin A. One $\mu\text{g/ml}$ of trichostatin A (manufactured by Wako Pure Chemical Industries, Ltd.) transactivates p51 in the cultured human cell line, and it was confirmed under this condition that the promoter activity of the gene having the isolated p51 promoter region was further remarkably enhanced. Thus, these results demonstrate that the gene having the base sequence as set forth in SEQ ID NO: 1 of the sequence listing has a function as a p51 promoter.

(1-2) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 1 of the sequence listing, and that has p51 promoter activity, is also encompassed in the scope of the present invention. The deletion, substitution, or addition of bases is to a degree that does not substantially affect the structure and function of the entire

promoter. The degree of deletion, substitution, or addition of these bases is acceptable when it has p51 promoter activity and it has a homology with the original base sequence of 80% or higher, preferably 90% or higher, and more preferably 99% or higher.

(1-3) DNA that hybridizes to DNA comprising the base sequence as set forth in SEQ ID NO: 1 of the sequence listing under a stringent condition and that has a p51 promoter activity is also encompassed in the scope of the present invention. Such hybridizing DNA mutants include DNA sequences partially altered by the mutation, deletion, ligation etc. of DNA fragments by site-directed mutagenesis, mutagen-treated random mutation, cleavage with restriction enzymes. The degree to which these DNA mutants hybridize to the coding gene as set forth in SEQ ID NO: 1 is a stringent condition: for example, the above membrane is incubated in a hybridization solution (50 mM Tris-HCl, pH 7.5, 1 M sodium chloride, 1% sodium dodecylsulfate, 10% dextran sulfate, 0.2 mg/ml yeast RNA, 0.2 mg/ml salmon sperm DNA) at 65°C for one hour as prehybridization, then a radioisotope-labeled cDNA fragment is added to one million dpm/ml in terms of the amount of radioisotope and incubated at 65°C for 16 hours as hybridization, and subsequently the membrane is washed in a 2 x SSC solution (300 mM sodium chloride, 30 mM trisodium citrate) containing 0.1% sodium dodecylsulfate at 65°C for 30 minutes, followed by autoradiography analysis to

confirm hybridization on an X-ray film.

(1-4) DNA that is shown in SEQ ID NO: 2 of the sequence listing and that encodes the p51 promoter region and the 5'-untranslated region of p51 is also encompassed in the scope of the present invention. The base sequence of positions from 5677 to 5960 in the SEQ ID NO: 2 corresponds to DNA encoding 5'-untranslated region. Among them, the base sequence of positions from 5767 to 5960 corresponds to the intron region.

(1-5) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing and that has p51 promoter activity is also encompassed in the scope of the present invention. The deletion, substitution, or addition of bases is to a degree that does not substantially affect the structure and function of the entire promoter. The degree of deletion, substitution, or addition of these bases is acceptable when it has p51 promoter activity and it has a homology with the original base sequence of 80% or higher, preferably 90% or higher, and more preferably 99% or higher.

(1-6) DNA that hybridizes to DNA comprising the base sequence as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition and that has p51 promoter activity is also encompassed in the scope of the present invention. Such hybridizing DNA mutants include those similar to those exhibited in the

above (1-3). The degree to which these DNA mutants hybridize to the gene as set forth in SEQ ID NO: 2 is such that hybridization is confirmed under a stringent condition similar to the one described in the above (1-3).

(2) Gene encoding the 5'-untranslated region of p51 of the present invention

(2-1) DNA that has a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing is DNA that encodes the 5'-untranslated region of p51 located downstream of the p51 promoter region, and such DNA is also encompassed in the scope of the present invention. Among them, the base sequence of positions from 5767 to 5960 corresponds to the intron region. Part or all of DNA that encodes the 5'-untranslated region of p51 may be used as a nucleic acid probe, for example, for cloning a gene as described below.

(2-2) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that has a function similar to that of the DNA in the above (2-1) is also encompassed in the scope of the present invention. The deletion, substitution, or addition of bases is to a degree that it does not substantially affect the structure and function of the DNA. The

degree of deletion, substitution, or addition of these bases is acceptable when it has a function similar to DNA described in the above (2-1) and it has a homology with the original base sequence of 80% or higher, preferably 90% or higher, and more preferably 99% or higher.

(2-3) DNA that hybridizes to DNA comprising the base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition, and that has a function similar to that of DNA in the above (2-1) is also encompassed in the scope of the present invention. Such hybridizing DNA mutants include those similar to those exhibited in the above (1-3). The degree to which these DNA mutants hybridize to the gene as set forth in SEQ ID NO: 2 is such that hybridization is confirmed under a stringent condition similar to the one described in the above (1-3).

(3) A recombinant plasmid containing a gene having the p51 promoter region of the present invention

By constructing a recombinant plasmid containing a gene having the p51 promoter region of the present invention, it is possible to allow Escherichia coli etc. to stably retain said gene, whereupon vectors used include any commonly used ones such as pBluescript II SK(-). In the examples described below, there is exhibited the pBS/p51 promoter in which a gene having the p51 promoter region has been integrated into

pBluescript II SK(-). After cleaving these plasmids with an appropriate restriction enzyme as needed, they can be ligated to an suitable vector to make plasmids for determining promoter activity. As a plasmid for
5 determining promoter activity, a plasmid such as pGL2 can be used as a vector.

(4) A transformant or a transductant

The above recombinant plasmid can be introduced into an appropriate host to construct a
10 transformant or a transductant. Escherichia coli, yeast, and mammalian cells can be used. A transformant that retains a plasmid for determining promoter activity can be obtained by transforming into an appropriate host a recombinant plasmid that has been
15 integrated into a vector for determining activity as described above. For example, a recombinant plasmid as shown in Figure 1 may be introduced into a cultured mamallian cell to obtain a transformant. A trans-
formant or a transductant may be cultured in an
20 appropriate nutrient culture medium and the amount of the reporter gene expressed in the cell may be measured to determine the promoter activity.

(5) A nucleic acid probe, and gene cloning and detection using it.

25 In accordance with the present invention, part or all of the gene having the p51 promoter region of the present invention or part or all of the gene encoding the 5'-untranslated region of p51 of the

present invention may be used as a nucleic acid probe for use in gene cloning etc. described below. As a nucleic acid probe comprising part of the gene of the present invention, there may be mentioned a nucleic acid probe comprising an oligonucleotide of 15 nucleotides or more. Said nucleic acid probe may be prepared by ligating the above gene of the present invention or a gene fragment thereof to a suitable vector, which is then introduced into a bacterium, replicated, extracted from the debris of cell homogenate with phenol etc., cleaving with the restriction enzyme that recognizes the site at which the vector is ligated, electrophoresing, and excising the probe from the gel. Said nucleic acid probe may also be prepared, based on the base sequence as set forth in SEQ ID NO: 1 or 2 of the sequence listing, by chemical synthesis using a DNA synthesizer or by a gene amplification technology with polymerase chain reaction (PCR). Said nucleic acid probe may be labeled with a radioisotope or fluorescence to enhance detection sensitivity at the time of use.

Said nucleic acid probe may be used in the cloning method of genes other than the above having a promoter activity or the cloning method of the protein-encoding genes located downstream thereof. Thus, by searching the genome library of various biological tissues by the hybridization method or the PCR method using said nucleic acid probe, it is possible to

isolate genes having a similar function to the gene of the present invention or protein-encoding genes located downstream thereof.

(6) Antisense DNA and antisense RNA

5 In accordance with the present invention, there may also be provided antisense DNAs and antisense RNAs of genes having the above-mentioned p51 promoter region as well as antisense DNAs and antisense RNAs of genes having the 5'-untranslated region of p51. By
10 introducing these antisense DNAs and antisense RNAs into a cell, the expression of genes encoding p51 can be suppressed or enhanced. As the antisense DNA to be introduced, there can be used, for example, an antisense DNA or part thereof corresponding to SEQ ID
15 NO: 1 or 2 of the sequence listing. An example of said antisense DNA is shown in SEQ ID NO: 3 of the sequence listing. This represents the sequence of the antisense DNA of a gene having the p51 promoter activity of SEQ ID NO: 1 of the sequence listing. As antisense DNA,
20 for example, there can be used a fragment obtained by suitably cleaving part of these antisense DNAs or DNAs synthesized based on the DNA sequences of these antisense DNAs.

 As antisense RNA, for example, there can be
25 used antisense RNA or part thereof corresponding to SEQ ID NO: 1 or 2 of the sequence listing. An example of said antisense RNA is shown in SEQ ID NO: 4 of the sequence listing. This represents the sequence of the

antisense RNA of the gene having the p51 promoter activity of SEQ ID NO: 1 of the sequence listing. As antisense RNA, for example, there can be used a fragment obtained by suitably cleaving part of these antisense RNAs or RNAs synthesized based on the RNA sequences of these antisense RNAs. A gene SEQ ID NO: 1 of the sequence listing having the p51 promoter activity or part thereof may be ligated to an appropriate vector, which is then introduced into a bacterium, replicated, extracted from the debris of cell homogenate with phenol etc., which is used as a template to allow RNA polymerase to act in an in vitro transcription system to synthesize RNA to be used. Antisense DNAs and antisense RNAs may be chemically modified so that they are refractory to in vivo decomposition and they can pass through cell membrane. The antisense DNAs and antisense RNAs thus prepared can be used for the treatment of various diseases including malignant tumor.

(7) Drug screening

In accordance with the present invention, there is provided a screening method for novel drugs using the transformant or the transductant of the present invention. For example, an agent that has the p51 promoter region activation activity and that enhances the expression of p51 protein can be discovered by screening drugs that enhance luciferase activity using a cultured cell that has integrated a

recombinant plasmid in which the luciferase reporter gene and said gene have been ligated.

As cultured cells into which a recombinant plasmid is to be introduced, any cultured cell lines that can be passaged can be used including a cultured colon cancer cell line HCT-116. As recombinant plasmids, any vector can be used that contains a reporter gene including pGL2-neo vector in which the neomycin gene has been ligated to the pGL2 plasmid used in Examples. In the construction of a screening system, a cultured cell line is selected into which a vector for screening is introduced by the lipofectin method, etc. By culturing this transformed cell together with the selection drug, cells that contain the vector for screening can only be grown. These cells may be used as they are in screening or may be cloned to be prepared as a single cell line and then used in screening. By adding a sample to these cells and then determining the activity of the reporter protein, agents that regulate the transcription of p51 can be searched. As samples, for example, microbial secondary metabolites may be used, or synthetic compounds may be used. Agents that activates the p51 promoter enhance the production of p51 protein and exhibit a growth suppressing effect on tumor cells, even if p53 is mutated in these cells, and thereby are expected to become novel anti-cancer agents.

(8) Pharmaceutical preparation

The p51's expression is only observed in highly localized tissues such as muscle cells. Thus, it can be used as a gene therapy vector by ligating a gene of interest downstream of the gene having the p51 promoter activity of the present invention, and allowing it to express the gene of interest only in specific tissues. The above antisense DNAs and antisense RNAs can be therapeutic agents for various disease including malignant tumor, also.

Now, the present invention will be specifically explained below. It is to be noted, however, that the present invention is not limited by these examples in any way.

Example 1

Isolation of a novel gene fragment encoding the p51 promoter region

(1-1) Preparation of a probe for screening

A probe for screening a novel gene fragment that encodes the p51 promoter region was prepared. As such a method, the RT-PCR method was used. The experimental procedure is shown below. From a six μ g of human muscle-derived RNA (manufactured by OriGene), cDNA was prepared using 200 units of a reverse transcriptase Superscript (manufactured by GIBCO BRL). With this as the template, the amplification of a p51 mRNA-derived gene fragment was attempted by a PCR

method using a primer set shown in SEQ ID NO: 5 and 6 and a primer set shown in SEQ ID NO: 7 and 8. The gene amplified using the former primer set encodes 143 bases of the 5'-untranslated region of p51A mRNA reported until now and 165 bases of the 5'-terminal end of the open reading frame, and the gene amplified using the latter primer set encodes the full-length of the p51A mRNA open reading frame. The base sequence of the former gene is shown in SEQ ID NO: 9 of the sequence listing. In performing the PCR method, a DNA polymerase EX Taq (manufactured by Takara Shuzo) was used, and the PCR condition comprised 30 cycles of amplification with each cycle comprising 95°C for one minute, 55°C for one minute, and 72°C for one minute. PCR amplified products were subjected to phenol treatment and chloroform treatment to remove protein, and then to ethanol precipitation. Then this DNA was washed in 70% ethanol and dissolved in a sterile water. After these purified DNAs and 20 ng of E. coli vector pBluescript KS(+) (manufactured by Toyobo) which was cleaved with a restriction enzyme EcoRV (manufactured by Takara Shuzo), and in which one thymine was added (referred to hereinafter as pBS/EcoRV TA), were ligated using a DNA ligation kit (manufactured by Takara Shuzo) (the plasmid containing the amplified gene obtained with the primer set shown in SEQ ID NO: 5 and 6 is hereinafter referred to as pBS/p51-1, and the gene containing the amplified gene obtained with the primer

set shown in SEQ ID NO: 7 and 8 is hereinafter referred to as pBS/p51A ORF). The absence of mutation in the amplified genes was confirmed by determining the base sequences of pBS/p51-1 and pBS/p51A ORF. The determi-

5 nation of the base sequence was performed using an automated sequencer LONG READIR4200 (manufactured by LICOR). Subsequently, pBS/p51-1 was cleaved with restriction enzymes EcoRV and PstI (manufactured by Toyobo), which were then fractionated on a 0.8% agarose
10 electrophoresis, extracted, and purified. For purification, EASYTRAP (manufactured by Takara Shuzo) was used. This purified DNA was used as a template for the screening probe.

(1-2) Screening of a novel gene fragment encoding the
15 p51 promoter region

Subsequently, the screening of gene fragments of a gene encoding the p51 promoter region from a human genomic library was attempted. As the human genomic library, Easy-to-Handle Eukaryotic Genomic Library from
20 human (manufactured by Mo Bi Tec) was used. The infection efficiency of this library was three million plaque forming units per microliter. Using this library, a membrane for plaque hybridization was prepared in the following manner. 0.02 μ l of the
25 library solution and 0.9 ml of C600 E. coli solution (E. coli strain C600 was cultured under shaking in 50 ml of the LB medium (0.5% yeast extract, 1% peptone, 0.5% sodium chloride) containing 0.2% maltose and 10 mM

magnesium sulfate) at 37°C for 16 hours, centrifuged at 5000 rpm for 5 minutes, recovered, and then suspended in 25 ml of 10 mM magnesium sulfate) were mixed, and incubated at 37°C for 15 minutes so as to be infected with phage. Seven ml of LB soft agar (0.7% agar was added to 0.5% yeast extract, 1% peptone, 0.5% sodium chloride and 10 mM magnesium sulfate to prepare an agar medium) that was incubated at 47°C after melting was added thereto, gently mixed, and then was plated onto a LB-magnesium sulfate plate (1.5% agar was added to 0.5% yeast extract, 1% peptone, 0.5% sodium chloride and 10 mM magnesium sulfate to prepare an agar medium) with an internal diameter of 150 mm. By this procedure, 60,000 plaques appeared per plate. A similar procedure was used to plate on a total of 23 plates. They were incubated at 37°C for 16 hours to form plaques. After the plate was cooled at 4°C for one hour, the plaques were adhered to a nylon membrane, Colony plaque screen (manufactured by NEN). After this membrane was dried, it was allowed to denature in 500 ml of an alkaline solution (0.2 N sodium hydroxide, 1.5 M sodium chloride) at room temperature for 2 minutes followed by neutralization in 500 ml of a neutralizing solution (0.5 M Tris-HCl, pH 7.2, 1.5 M sodium chloride, 1 mM ethylenediaminetetraacetic acid disodium salt) at room temperature for 2 minutes. This membrane was then incubated in a 3 X SSC solution (450 mM sodium chloride, 45 mM trisodium citrate) at 55°C for one

hour, and dried to prepare a membrane for plaque hybridization. Fifty ng of the above purified cDNA fragment obtained by cleaving pBS/p51-1 with restriction enzymes EcoRV and PstI was labeled using Prime-It II (manufactured by Stratagene) to prepare a probe for screening.

Then the membrane was used for plaque hybridization. The above membrane was incubated in 50 ml of a hybridization solution (50 mM Tris-HCl, pH 7.5, 1 M sodium chloride, 1% sodium dodecylsulfate, 10% dextran sulfate, 0.2 mg/ml yeast RNA, 0.2 mg/ml salmon sperm DNA) at 65°C for one hour as prehybridization. Then a radioisotope-labeled cDNA fragment was added to one million dpm/ml in terms of the amount of radioisotope and incubated at 65°C for 16 hours as hybridization. Subsequently the membrane was washed in a 2 x SSC solution (300 mM sodium chloride, 30 mM trisodium citrate) containing 0.1% sodium dodecylsulfate at 65°C for 30 minutes. The washing was performed twice. Then, the membrane was subjected to autoradiography to detect positive plaques.

Then, plaque groups containing plaques corresponding to positive signal were isolated, and suspended in one ml of the SM buffer solution (50 mM Tris-HCl, pH 7.5, 100 mM sodium chloride, 10 mM magnesium sulfate, 0.01% gelatin). The solution was incubated at 4°C for 16 hours to elute phage, and the phage was recovered in the supernatant fluid after

centrifuging at 13000 rpm for 10 minutes. The phage solution was plated on a LB-magnesium sulfate plate in a manner similar to the above to screen positive plaques. As a result, positive plaques were successfully isolated as completely isolated plaques. From these plaques, phage was isolated in a manner similar to the above.

Subsequently, the following experiment was performed in order to determine the base sequence of the library DNA contained in this phage. The library has been made using lambda PS phage (manufactured by Mo Bi Tec). The phage has the recombinant site of loxP in the base sequence thereof, and thereby it can be introduced into an E. coli strain BNN132 having Cre recombinase in order to excise the vector portion containing the genome library sequence from the phage. This procedure was carried out as described below. Twenty μ l of the phage solution was mixed with 200 μ l of the BNN132 E. coli solution (E. coli strain BNN132 was cultured under shaking in 50 ml of the LB medium containing 0.2% maltose and 10 mM magnesium sulfate at 37°C for 16 hours, and then recovered by centrifuging at 5000 rpm for 5 minutes, which was then suspended in 25 ml solution of 10 mM magnesium sulfate) and incubated at 37°C for 30 minutes. The mixture was then plated onto the LB-ampicillin plate (1.5% agar was added to 0.5% yeast extract, 1% peptone, 0.5% sodium chloride and 0.1 mg/ml ampicillin sodium to prepare an

agar medium), and the plasmid of interest was prepared from the colonies appeared. The preparation of the plasmid was performed according to the procedure described in Lab Manual Gene Engineering (Published by Maruzen, edited by Masami Muramatsu, 1990) on pages 53-55. The plasmid was found to contain about 15 kb of the library gene. This is referred to hereinafter as pPS/library.

(1-3) Sequencing of the gene fragment encoding the p51 promoter region

Then, part of the base sequence of the library gene in pPS/library was determined using an automated sequencer LONG READIR4200 to confirm that it contains part of the cDNA sequence complementing p51A mRNA. It was also found that a 0.6 kb gene fragment obtained by cleaving of pPS/library with a restriction enzyme EcoRV contains part of the cDNA sequence that complements p51A mRNA. The pPS/library was then cleaved with a restriction enzyme PvuII (manufactured by Toyobo), fractionated on 0.8% agarose electrophoresis, and blotted on a nylon membrane Hybond N+ (manufactured by Amersham). Blotting of the plasmid fragment was performed according to the procedure described in Lab Manual Human Genome Mapping (published by Maruzen, edited by Masaaki Hori and Yusuke Nakamura, 1991) on pages 26-36. This membrane was used in Southern blotting according to the following procedure. As a cDNA fragment for probing, pPS/library

was used and cleaved with a restriction enzyme EcoRV,
and after fractionation on a 0.8% agarose electro-
phoresis a 0.6 kb gene fragment was excised and
purified using EASYTRAP. This purified cDNA was
5 radiolabeled using Prime-It II to prepare a probe. The
membrane was incubated in a hybridization solution for
Southern blotting (10% sodium dodecylsulfate, 7%
PEG8000) at 65°C for one hour as prehybridization.
Then the above probe was added to one million dpm/ml in
10 terms of the amount of radioisotope and incubated at
65°C for 16 hours for hybridization. Subsequently, the
membrane was washed in a 2 x SSC solution containing
0.1% sodium dodecylsulfate at 65°C for 30 minutes. The
washing was performed twice. Then, the membrane was
15 subjected to autoradiography to demonstrate that a
PvuII-cleaved fragment containing the immediate
upstream region of p51A mRNA was 5.5 kb in length.
Then, pPS/library was cleaved with a restriction enzyme
PvuII, and after fractionation on a 0.8% agarose
20 electrophoresis a 5.5 kb gene fragment was excised and
purified using EASYTRAP. The purified cDNA fragment
was ligated, using the DNA ligation kit, to pBluescript
II Sk(+) that had been cleaved with a restriction
enzyme EcoRV followed by dephosphorylation with shrimp
25 alkaline phosphatase. The plasmid is referred to
hereinafter as pBS/PvuII5.5. Its base sequence was
determined using the automated sequencer LONG READIR
4200. The base sequence and the partial base sequence

of the pPS/library determined above were combined to obtain the base sequence as set forth in SEQ ID NO: 2 of the sequence listing. Furthermore, a base sequence obtained by removing the part that encodes the 5'-

5 untranslated region of p51 from the base sequence as set forth in SEQ ID NO: 2 of the sequence listing is shown in SEQ ID NO: 1 of the sequence listing. The 3'-terminal of this PvuII-cleaved fragment was found to be located about 0.22 kb upstream of the base at position
10 1 of SEQ ID NO: 9 of the sequence listing that is a cDNA sequence corresponding to p51A mRNA.

Accordingly, in order to prepare a plasmid containing bases up to the base at position 1 of SEQ ID NO: 9 of the sequence listing that is cDNA correspond-
15 ing to part of p51A mRNA, the following procedure was performed. The pPS/library was cleaved with a restriction enzyme EcoRV, which was fractionated on a 0.8% agarose electrophoresis and then a 0.6 kb gene fragment was excised and purified using EASYTRAP, which was
20 ligated using a DNA ligation kit to pBS/PvuII5.5 that had been cleaved with a restriction enzyme EcoRV and a restriction enzyme SmaI (manufactured by Takara Shuzo) followed by dephosphorylation with shrimp alkaline phosphatase. The plasmid obtained was analyzed and the
25 plasmid that was ligated in the correct direction was isolated. The plasmid is referred to hereinafter as pBS/p51 promoter.

In order to demonstrate that the base

sequence is surely present on the genome, the following experiment was performed. From a colon cancer cell line HCT116, genomic DNA was prepared, which was used as a template in a PCR. The genomic DNA was prepared according to Cell Engineering Experimental Protocol (published by Shujunsha, edited by Tokyo University, the Institute of Medical Science, Department of Anti-Cancer Research, 1993) on pages 16-19. The primers used are those shown in SEQ ID NO: 10 and 11. Respectively, they correspond to the sense strand of the bases at positions from 3543 to 3570 and the antisense strand of the bases at positions from 5458 to 5487 of the base sequence as set forth in SEQ ID NO: 1 of the sequence listing. Polymerase used was LA Taq (manufactured by Takara Shuzo), and PCR comprised 30 cycles of amplification with each cycle comprising 94°C for one minute and 68°C for three minutes. The PCR products were subjected to a 0.8% agarose electrophoresis to confirm that a gene fragment of 1.9 kb was specifically amplified. Furthermore, after the fragment was excised, it was purified using EASYTRAP, and ligated to pBS/EcoRV TA using a DNA ligation kit. The base sequence of this plasmid was determined using an automated sequencer LONG READIR 4200, and it was confirmed to completely match the base sequence in the pBS/p51 promoter.

(1-4) Construction of a luciferase vector containing the p51 promoter region

In order to confirm that promoter activity is present on the isolated region, a vector in which the gene sequence was ligated upstream of the luciferase reporter gene was prepared according to the following method. As the vector having the luciferase reporter gene, pGL2 (manufactured by Promega) was used. First, to allow for integration of pGL2 and selection in mammalian cells, and a neomycin-resistant gene was introduced into this vector. The pGL2 cleaved with a restriction enzyme BamHI (manufactured by Toyobo) was ligated using a DNA ligation kit to a 2617-base neomycin-resistant gene, being obtained by cleaving pMAM neo (manufactured by Toyobo) with BamHI and fractionating on a 0.8% agarose gel electrophoresis excised and purified using EASYTRAP. The plasmid is referred to hereinafter as pGL2-neo. Subsequently, pGL2-neo was cleaved with a restriction enzyme XhoI (manufactured by Toyobo), precipitated with ethanol, washed in 70% ethanol, and dissolved in purified water. The cleaved end of this DNA was blunt-ended using a DNA blunting kit (manufactured by Takara Shuzo), precipitated with ethanol, washed in 70% ethanol, and dissolved in purified water. Then, this DNA was cleaved with a restriction enzyme KpnI (manufactured by Boehringer Mannheim), treated with phenol, treated with chloroform to remove protein, precipitated with

ethanol, washed in 70% ethanol, and dissolved in sterile water. This DNA is referred to hereinafter as pGL2-neo/XhoI (blunting), KpnI.

Subsequently, the pBS/p51 promoter was

5 cleaved with a restriction enzyme NotI (manufactured by Boehringer Mannheim), precipitated with ethanol, washed in 70% ethanol, and dissolved in purified water. After the cleaved end of this DNA was blunt-ended using a DNA blunting kit (manufactured by Takara Shuzo), precipi-
10 tated with ethanol, washed in 70% ethanol, and dissolved in purified water. Then, this DNA was cleaved with a restriction enzyme KpnI (manufactured by Boehringer Mannheim), fractionated on a 0.8% agarose electrophoresis, and a 5.7 kb gene fragment was excised
15 and purified using EASYTRAP. This gene fragment is referred to hereinafter as pBS/p51 promoter/NotI (blunting), KpnI5.7. Then, pGL2-neo/XhoI (blunting), KpnI and pBS/p51 promoter/NotI (blunting), KpnI5.7 were ligated using a DNA ligation kit. This plasmid is
20 referred to hereinafter as pGL2-neo/p51 promoter. The gene sequence of this plasmid is shown in SEQ ID NO: 12 of the sequence listing. The restriction map of this plasmid is shown in Figure 1.

(1-5) Effect of various drugs on the amount of p51
25 transcribed

1) Northern hybridization

A variety of stimulations including stimulation with a drug were applied to a cell line HCT116

derived from colon cancer to investigate changes in the
 amount expressed of p51 by Northern blotting. 200,000
 cells of HCT116 cells were plated on a culture dish
 with a diameter of 60 mm, and cultured in the presence
 5 of 5% carbon dioxide at 37°C for two days. The medium
 used was McCOY's 5A (manufactured by Nisseiken) to
 which fetal bovine serum (FBS) was added to a final
 concentration of 10% (referred to hereinafter as
 McCOY's 5A/10% FBS). Then it was treated with A23187
 10 (0.1 μ M, 1 μ M), cisplatin (1 μ g/ml, 10 μ g/ml), CPTcAMP
 (0.1 μ M, 1 μ M), etoposide (10 ng/ml, 0.1 μ g/ml),
 genistein (1 μ M, 10 μ M), ML-236B (1 μ g/ml, 10 μ g/ml),
 milurinine (1 μ M, 10 μ M), mitomycin C (5 μ g/ml, 50
 μ g/ml), NKH477 (0.1 μ M, 1 μ M), okadaic acid (1 nM, 10
 15 nM), radicicol (0.1 μ g/ml, 1 μ g/ml), staurosporin (1
 nM, 10 nM), taxol (1 μ g/ml, 10 μ g/ml), trichostatin A
 (0.1 μ g/ml, 1 μ g/ml), radiation (50 J/m², 100 J/m²),
 vitamin D₃ (10 nM, 100 nM), vincristine (1 μ g/ml, 10
 μ g/ml), and wortmannin (0.1 μ M, 1 μ M), and further
 20 cultured in the presence of 5% carbon dioxide at 37°C
 for 24 hours. Then, from the cells, RNA was prepared
 using ISOGEN (manufactured by Nippongene). 20 μ g each
 of RNA was fractionated on a 1% agarose electrophoresis
 containing formaldehyde, and blotted on a nylon
 25 membrane HybondN+ according to the method in the above
 New Cell Engineering Experimental Protocol on pages
 194-197. As the probe, pBS/p51A ORF was cleaved with a
 restriction enzyme EcoRI (manufactured by Toyobo),

fractionated on a 0.8% agarose electrophoresis, and then a 0.6 kb gene fragment was excised and purified using EASYTRAP, which was radiolabeled with Prime-It II and used. The membrane was incubated in 10 ml of a prehybridization solution (25 mM phosphate buffer solution, pH 7.0, 6 X SSC, 50% formamide, 5 X Denhardt's solution, 0.1% sodium dodecylsulfate, 0.2 mg/ml salmon sperm DNA) at 42°C for four hours as prehybridization. Then it was replaced with 10 ml of a hybridization solution (20 mM phosphate buffer solution, pH 7.0, 6 X SSC, 50% formamide, 1 X Denhardt's solution, 0.1% sodium dodecylsulfate, 0.1 mg/ml salmon sperm DNA, 4% dextran sulfate), and then the above radiolabeled probe was added to one million dpm/ml and incubated at 42°C for 16 hours as hybridization. Subsequently, it was washed in a 2 x SSC solution containing 0.1% sodium dodecylsulfate at 65°C for 30 minutes. The washing was performed twice. Then, the membrane was analyzed on autoradiography.

2) Result

As a result, a specific band that reacts with p51A-derived cDNA was only observed in the cells treated with 1 µg/ml trichostatin A. In any other stimulation, no increases in the amount of p51A mRNA were observed. Figure 2 shows an increase in the amount of p51A mRNA with trichostatin A treatment. Figure 2 shows blotting of 20 µg of RNA from: lane 1 and 2: cells treated with taxol (1 µg/ml, 10 µg/ml),

lane 3 and 4: cells treated with trichostatin A (0.1 $\mu\text{g/ml}$, 1 $\mu\text{g/ml}$), lane 5 and 6: cells treated with radiation (50 J/m^2 , 100 J/m^2), lane 7 and 8: cells treated with vitamin D₃ (10 nM, 100 nM), lane 9 and 10: cells treated with vincristine (1 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$), and lane 11 and 12: cells treated with wortmannin (0.1 μM , 1 μM).

(1-6) Functional analysis of the p51 promoter region

Subsequently, the following experiment was carried out in order to perform functional analysis of the p51 promoter region. 200,000 cells of HCT116 were plated on a culture dish with a diameter of 60 mm. The medium used was McCoy's 5A/10% FBS. These were cultured in the presence of 5% carbon dioxide at 37°C for 48 hours. The medium was replaced with McCoy's 5A containing no serum. Subsequently, 2 μg each of pGL2-neo/p51 promoter or pGL2-neo vector was introduced. The plasmid was introduced using the lipofectamine plus reagent (manufactured by GIBCO BRL). Thus, 125 μl of McCoy's 5A and 8 μl of the plus solution were added to the plasmid solution, stirred well, and then incubated at room temperature for 15 minutes. To this solution, 125 μl of McCoy's 5A and 12 μl of lipofectamine solution were added, stirred well, and then incubated at room temperature for further 15 minutes. This was added to the HCT116 cells incubated by the above procedure, gently mixed, and cultured in the presence of 5% carbon dioxide at 37°C for three hours to

introduce plasmid. Subsequently, the medium was replaced with McCoy's 5A/10% FBS, and cultured in the presence of 5% carbon dioxide at 37°C for 21 hours.

Then, trichostatin A was added to two groups of cells to a final concentration of 0 or 1 µg/ml, and cultured in the presence of 5% carbon dioxide at 37°C for 24 hours. For these cells, the luciferase activity was determined to investigate promoter activity. As a negative control, cells containing no luciferase gene, pcDNA3/lacZ (the lacZ gene ligated to the mammalian cell expression vector pcDNA3) was transfected in a procedure similar to the above. The luciferase activity was determined according to the following method.

The medium in the cell culture dish was removed, and 5 ml of the cell lysis reagent (25 mM Tris-HCl, pH 7.8, 2 mM dithiothreitol, 2 mM CDTA, 0.2% triton X-100, 10% glycerol) was added and incubated at room temperature for 15 minutes. The cell lysis solution was taken at 10, 20, 30, 40, and 50 µl aliquotes, to which 50 µl of the substrate solution (20 mM Tricine-sodium hydroxide, pH 7.8, 1.07 mM basic magnesium carbonate, 2.67 mM magnesium sulfate, 0.1 mM ethylenediaminetetraacetic acid disodium salt, 33.3 mM dithiothreitol, 270 µM coenzyme A, 470 µM luciferine, 530 µM adenosine triphosphate) was added, and incubated at room temperature for 15 seconds, and its chemiluminescence intensity was determined using a

luminometer LUMINOSKAN (manufactured by Dainippon Pharmaceutical). The result is shown in Figure 3.

These results demonstrated that pGL2-neo/p51 promoter-introduced cell lysis solution exhibits a significantly high fluorescence intensity compared to pGL2-neo. The administration of trichostatin A also markedly enhanced fluorescence intensity in the pGL2-neo/p51 promoter. The result is in agreement with the above result in Figure 2.

The above experimental data demonstrated that the isolated gene fragment is a functional p51 promoter region.

Example 2

Screening of drugs that modify the p51 promoter function

(2-1) Preparation of a cultured cell line for screening

The cultured cell line retaining the pGL2-neo/p51 promoter plasmid can be used as cells for screening drugs that modify the p51 promoter function.

Said drug is effective for treatment of various diseases including cancer. Said cells were prepared according to the following procedure.

As the plasmid-introduced cells, the HCT116 cell line was used. 500,000 cells of HCT116 cell were plated on a culture dish with a diameter of 60 mm. The medium used was McCoy's 5A/10% FBS. These were cultured in the presence of 5% carbon dioxide at 37°C

for 48 hours. The medium was replaced with McCoy's 5A containing no serum. As the plasmid for introduction, two μg of pGL2-neo/p51 promoter was dissolved in 4 μl of the TE solution to prepare a plasmid solution. The
5 plasmid was introduced using the lipofectamine plus reagent (manufactured by GIBCO BRL). Thus, 125 μl of McCoy's 5A and 8 μl of the plus solution were added to the plasmid solution, stirred well, and then incubated at room temperature for 15 minutes. To this solution,
10 125 μl of McCoy's 5A and 12 μl of the lipofectamine solution were added, stirred well, and then incubated at room temperature for further 15 minutes. This was added to the HCT116 cells cultured by the above procedure, gently mixed, and incubated in the presence
15 of 5% carbon dioxide at 37°C for three hours to permit the introduction of the plasmid. Subsequently, the medium was replaced with McCoy's 5A/10% FBS, and incubated in the presence of 5% carbon dioxide at 37°C for 21 hours. Then, the cells were recovered from the
20 culture dish with PBS containing 0.25% trypsin and 0.02% EDTA, and 1% amount thereof was plated onto a culture dish with a diameter of 100 mm. It was further cultured in McCoy's 5A/10% FBS medium in the presence of 5% carbon dioxide at 37°C for further 72 hours, and
25 then 1.2 mg/ml of geneticin (manufactured by Nacalai Tesque) was added as a selection drug. By replacing the medium every three days, and further culturing in the above condition for further two weeks, the cells

into whose genes the plasmid gene was introduced, and became geneticin-resistant, were selected.

Subsequently, 50 clones were isolated from the cell group that formed colonies on the dish, and were plated
5 on a 24-well culture dish to continue being cultured.

These clones are expected to exhibit an enhanced luciferase activity to stimulation such as the activation of the p51 promoter function. Thus, the reactivity of these clones to trichostatin A was
10 investigated by the following procedure. 500,000 cells each of the cloned cells were plated on a culture dish with a diameter of 60 mm. The medium used was McCoy's 5A/10% FBS. These were cultured in the presence of 5% carbon dioxide at 37°C for 48 hours. Then, tricho-
15 statin A was added to the two groups of cells to a final concentration of 0 or 0.1 µg/ml, and cultured in the presence of 5% carbon dioxide at 37°C for 24 hours. For these cells, their luciferase activity was determined to examine the effect on the p51 promoter
20 activity. The luciferase activity was determined according to the above method. As a result, several clones were obtained that exhibit a significantly enhanced luciferase activity to 0.1 µg/ml trichostatin A, from among which HCT116/p51 promoter clone #9 was
25 selected as the cell for screening. The result of luciferase assay is shown in Table 1.

Table 1

Clone No.	#5	#9	#17	#18
Without TsA	3.9	1.0	17.0	6.8
With TsA	29.5	33.3	44.1	14.0

In the Table, TsA represents trichostatin A. The trichostatin A treatment group received at 0.1 µg/ml. Values in the table show the measured values of luciferase activity.

5 (2-2) Screening of drugs that modify the p51 promoter function

HCT116/p51 promoter clone #9 cultured cell line is useful for screening drugs that induce the activation of the p51 promoter function. Said screening was performed according to the following procedure.

10 10,000 cells of HCT116/p51 promoter clone #9 cultured cell line were plated onto a 96-well culture dish, and cultured in 200 µl of RPMI1640/10% FBS in the presence of 5% carbon dioxide at 37°C for 24 hours.

15 Subsequently, a drug for screening was added, culturing was continued for 24 hours, and the luciferase activity was determined. As the drug, synthetic compounds or microbial secondary metabolites were used.

20 1200 compounds and 9600 microbial secondary metabolites were screened and 71 samples that is capable of activating the p51 promoter activity at a

level equal to or greater than that of 0.1 µg/ml trichostatin A. Substances that activate the p51 promoter are expected to become therapeutic agents for treating diseases including cancer.

5 INDUSTRIAL APPLICABILITY

In accordance with the present invention, there are provided a gene encoding the promoter region of protein p51 which is capable of inducing cell death, and a gene encoding the 5'-untranslated region of p51.

- 10 They are useful in diagnosing and treating diseases including cancer caused by abnormal regulation of cell propagation. Furthermore, in accordance with the present invention, there are provided the antisense DNAs and the antisense RNAs of these genes, nucleic
- 15 acid probes comprising parts or all of these genes, methods of detecting the above genes of the present invention or analogous genes using said nucleic acid probes, transformants in which the above gene of the present invention has been introduced, and methods of
- 20 screening drugs using them. These genes are also useful in diagnosing and treating diseases including cancer, etc.

SEQUENCE LISTING FREE TEXT

- The base sequences of SEQ ID NO: 5-8, 10 and
- 25 11 of the sequence listing show PCR primers.

The base sequence of SEQ ID NO: 9 of the

sequence listing shows the 5'-untranslated region of p51A mRNA and encodes 165 bases at the 5'-end of the open reading frame of p51A mRNA.

The base sequences of SEQ ID NO: 12 of the
5 sequence listing shows the DNA sequence of a plasmid containing the p51 promoter and the neomycin-resistant gene.

4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2

CLAIMS

1. A gene encoding the p51 promoter region shown in the following (1), (2), (3), (4), (5), or (6):

(1) DNA that encodes the p51 promoter region having the base sequence as set forth in SEQ ID NO: 1 of the sequence listing;

(2) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 1 of the sequence listing, and that has p51 promoter activity;

(3) DNA that hybridizes to the base sequence as set forth in SEQ ID NO: 1 of the sequence listing under a stringent condition, and that has p51 promoter activity;

(4) DNA that has the base sequence as set forth in SEQ ID NO: 2 of the sequence listing and that encodes the p51 promoter region and the 5'-untranslated region of p51;

(5) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that has p51 promoter activity; and

(6) DNA that hybridizes to the base sequence as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition, and that has p51 promoter activity.

2. A gene encoding the 5'-untranslated region of p51 shown in the following (7), (8), or (9):

(7) DNA that has a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing;

(8) DNA that has a base sequence in which one or a plurality of bases have been deleted, substituted, or added in a base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing, and that has a function similar to that of DNA in the above (7); and

(9) DNA that hybridizes to DNA comprising the base sequence of positions from 5677 to 5960 in the base sequence as set forth in SEQ ID NO: 2 of the sequence listing under a stringent condition, and that has a function similar to that of DNA in the above (7).

3. A recombinant plasmid comprising the gene of claim 1.

4. A transformant or a transductant comprising the recombinant plasmid of claim 3.


5. A nucleic acid probe comprising all or parts of the gene of claim 1 or 2.

6. A method of cloning the p51 promoter region or an analogous gene using the nucleic acid probe of claim 5.

7. A DNA sequence that is an antisense to all or parts of the gene of claim 1 or 2 and that enhances or inhibits the expression of p51 gene.


8. A RNA sequence that is an antisense to all or parts of the gene of claim 1 or 2 and that enhances or inhibits the expression of the p51 gene.
9. A method of screening drugs that act on p51 promoter using the transformant or the transductant of claim 4.
10. A compound that enhances or inhibits the expression of the p51 gene, said compound being selected by the screening method of claim 9.
11. A pharmaceutical formulation comprising the DNA sequence of claim 1, the DNA sequence of claim 7, or the RNA sequence of claim 8.

FIG. 1



2/3


FIG. 2



10/030294 2/3

3 / 3

FIG. 3



ES-95-01 #

COMBINED DECLARATION AND POWER OF ATTORNEY

(宣誓書及び委任状)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

"GENE ENCODING PROMOTER REGION OF TUMOR SUPPRESSOR GENE P51 AND USE THEREOF"

the specification of which: (check one) ☐ is attached hereto.

☒ was filed on June 28, 2000
as Application Serial No. PCT/JP00/04261
and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended, by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me which is material to patentability in accordance with Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date earlier than that of the application(s) on which priority is claimed:

Prior Foreign Application(s)			Priority Claimed	
<u>11-183195</u>	<u>Japan</u>	<u>June 29, 1999</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
_____	_____	_____	<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
_____	_____	_____	<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented, pending, abandoned)
_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented, pending, abandoned)

(Continued on Page 2)

ES295-01(7)

I hereby appoint the following as my attorneys of record with full power of substitution and revocation to prosecute this application and to transact all business in the Patent and Trademark Office:

③ Henry C. Nields, Reg. No. 17,029
Kevin S. Lemack, Reg. No. 32,579
Brian M. Dingman, Reg. No. 32,729
Please direct all communications to the following address:

Nields, Lemack & Dingman
176 E. Main Street - Suite 8
Westboro, Mass. 01581
U. S. A.

Telephone: (508) 898-1818

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

宣誓日

発明者フルネームサイン

氏名タイプ欄

-00 Date Dec. 12, 2001 Inventor Toshiyuki Sakai Toshiyuki SAKAI

Residence Kyoto-shi, Kyoto, Japan JPX Citizenship Japan

Post Office Address 28-1, Matsugasaki Kowakicho, Sakyo-ku, Kyoto-shi, Kyoto, Japan.

1-00 Date Dec. 12, 2001 Inventor Shigehide Kagaya Shigehide KAGAYA

Residence Omiya-shi, Saitama, Japan JPX Citizenship Japan

Post Office Address 2-336, Kitabukurocho, Omiya-shi, Saitama, Japan.

3-00 Date Dec. 12, 2001 Inventor Takamichi Sato Takamichi SATO

Residence Yono-shi, Saitama, Japan JPX Citizenship Japan

Post Office Address 6-8-22, Kamiochiai, Yono-shi, Saitama, Japan.

4-00 Date Dec. 12, 2001 Inventor Yoshikazu Sukenaga Yoshikazu SUKENAGA

Residence Tokyo, Japan JPX Citizenship Japan

Post Office Address 2-21-15-102, Ebaracho, Nakano-ku, Tokyo, Japan.

5-00 Date Dec. 12, 2001 Inventor Hideji Fujii Hideji FUJII

Residence Tokyo, Japan JPX Citizenship Japan

Post Office Address 3-48-21-1201, Honmachi, Shibuya-ku, Tokyo, Japan.

Date _____ Inventor _____

Residence _____ Citizenship _____

Post Office Address _____